

The British University in Egypt

BUE Scholar

Dentistry

Health Sciences

Winter 12-1-2023

Root and canal morphology of mandibular second molars in an Egyptian subpopulation: a cone-beam computed tomography study

Shehabeldin Mohamed Saber Prof.Dr.

The British University in Egypt, shehabeldin.saber@bue.edu.eg

Follow this and additional works at: <https://buescholar.bue.edu.eg/dentistry>



Part of the [Endodontics and Endodontology Commons](#)

Recommended Citation

Saber, Shehabeldin Mohamed Prof.Dr., "Root and canal morphology of mandibular second molars in an Egyptian subpopulation: a cone-beam computed tomography study" (2023). *Dentistry*. 197.
<https://buescholar.bue.edu.eg/dentistry/197>

This Article is brought to you for free and open access by the Health Sciences at BUE Scholar. It has been accepted for inclusion in Dentistry by an authorized administrator of BUE Scholar. For more information, please contact bue.scholar@gmail.com.

RESEARCH ARTICLE

Open Access



Root and canal morphology of mandibular second molars in an Egyptian subpopulation: a cone-beam computed tomography study

Shehabeldin Mohamed Saber^{1,2,3}, Mohammed abou El Seoud¹, Shaimaa Mohamed Abu el Sadat⁴ and Nawar Naguib Nawar^{1*} 

Abstract

Background This study was using Cone beam computed tomography (CBCT) to examine the anatomical variations in mandibular second molars in an Egyptian sub-population.

Methods A total of 350 CBCT images (215 females and 135 males, aged 15–65 years) of mandibular second molars were evaluated. Samples were evaluated in terms of: number of roots, number of root canals, roots' cross section as well as prevalence and configurations of C-shaped canals. Statistical analysis was done to highlight differences between different categories and their prevalence among genders (significance level was set at $p < 0.05$). Data were presented as frequency and percentage values and were analyzed using chi square test followed by pairwise comparisons utilizing multiple Fisher's exact tests with Bonferroni correction. The significance level was set at $p < 0.05$ within all tests.

Results Of the 350 mandibular second molars evaluated, 87.2% were non-C-shaped while 12.8% were C-shaped with no gender-based statistically significant differences ($\chi^2 = 0.19$, $p = 0.656$). Most samples had three root canals (80%) followed by two (16%), then one (3.2%), and finally four (0.8%) root canals. Among the non-C-shaped molars, presence of two roots was most common (83.4%) followed by presence of a single root (16.2%), and only one sample (0.2%) had three roots, and this had no correlation with gender ($\chi^2 = 1.86$, $p = 0.431$). In the mesial roots Type IV Vertucci was the most common configuration found (68.8%), while Type I was the most prevalent in the distal roots (91.8%). The long oval configuration was the most commonly found cross section in mesial roots while "oval" was the most prevalent in distal roots.

Conclusion Egyptian sub-population shows highly variable morphological features in mandibular second molars, hence, CBCT is highly recommended on case-to-case conditions.

Keywords C-shaped canal, Cone beam computed tomography, Endodontics, Mandibular second molar

Approved by Research Ethics Committee of The British University (El-Shorouk city, Egypt) (22-007).

*Correspondence:

Nawar Naguib Nawar
Nawar.Naguib@bue.edu.eg

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Sufficient knowledge on root canal morphology is essential for recognizing and treating anatomical details within the root canal system. Root canal systems show wide variations that include the presence of extra roots, extra root canals, fins, isthmuses loops, as well as unique morphologies such as S-shaped and C-shaped canals [1, 2]. Such anatomical details can shelter microbes, thus compromising the outcome of root canal treatment [1]. Hence, it is important to comprehend root canal morphologic variations in all tooth types before treatment [2].

Mandibular second molars are amongst the teeth that exhibit a wide variation in their internal and external anatomy according to race and geographic origin [3–14]. A characteristic anatomic variant that can exist in mandibular second molars is the C-shaped canal configuration, with a wide range of reported incidence (2.7–52%) [4, 15, 16]. It was first described by Cooke and Cox [17] and was attributed to the lack of fusion of Hertwig's root epithelial sheath of the vestibular or lingual side [17, 18]. It is characterized by a cross-sectional morphology of the pulp chamber that lacks the discrete orifices but rather resembles the letter "C", as long as the presence of a fin that connects separate root canals [4, 19]. Moreover, C-shaped root canals have thin lingual walls, thus adding to the challenges of different root canal treatment phases [20, 21]. Mandibular second molars might also have a supernumerary root; the radix entomolaris when it exists disto-lingually, or the radix paramolaris when located mesio-buccally [22].

Different methodologies evolved throughout decades to study the morphology of root canal systems [23]. One of the technologies that helped progress this process is the Cone beam computed tomography (CBCT) which is a non-invasive technology that allows a 1:1 accurate 3-dimensional evaluation of tooth dimensions, root and canal morphology [24, 25], and it was employed to examine various tooth types both experimentally and clinically [26–29].

To date, there is scarce knowledge about the morphology of mandibular second molars in the Egyptian population. This study aimed to investigate the number of roots, root canal configurations, prevalence and morphology of the C-shaped root canal system in mandibular second molars within an Egyptian subpopulation using high-resolution CBCT.

Methods

Sample size calculation and study design

A power analysis was performed using G*Power version 3.1.9.7. (Heinrich-Heine-Universität, Düsseldorf, Germany) based on the results of Senan et al. [30], the predicted sample size (n) was a total of (350) cases. The

study design, execution and reporting were planned to adhere to the "Preferred Reporting Items for Epidemiologic Cross-sectional Studies on Root and Root Canal Anatomy Using Cone-beam Computed Tomographic Technology" [31].

Subjects

This study was approved and the need for consent to participate was waived by the Research Ethics Committee of The British University (El-Shorouk city, Egypt) (22–007) given the type of the retrospective study and given that the CBCT images included were anonymous. Random CBCT images for 350 patients who required a preoperative 3D radiographic assessment between January 2021 and December 2021 were collected from a private maxillofacial imaging center and included in the study. The patients were referred to the imaging center for preoperative assessment as part of their dental examination, diagnosis, and treatment planning; this included orthodontic, periodontal and surgical reasons.

Inclusion and exclusion criteria

CBCT scans of patients who met the following criteria were included in the study: (i) Egyptian citizenship (ii) 15–65 years old (iii) presence of a fully mature and erupted mandibular second molar. Molars with root canal fillings, posts, crowns, internal or external root resorption, extensively calcified root canals, extensive coronal and root caries and periradicular radiolucency were excluded. The required sample size ($n=350$) was reached after examination of a total of 470 CBCT scans. The excluded scans ($n=120$ / 25% of total examined scans) did not fulfil the inclusion criteria.

Image acquisition

High resolution CBCT scans were acquired using Planmeca Promax 3D Classic unit (Planmeca, Helsinki, Finland) with exposure settings of 90 kV, 12.5 mA, 14.966 s, Field of view (FOV) 5.0×5.0 cm (one side of the mandible), and voxel size: 75 μm . Images' acquisition was done by an experienced radiologist. All data from the CBCT examinations were acquired in a digital DICOM format, imported to Planmeca Romexis dental imaging software (Planmeca, Helsinki, Finland), and viewed on an 18.5-inch HD LED monitor with a resolution of 1366 Å ~ 768.

Calibration and measurements

Examination was performed in March 2022 as detailed in many previous studies [16, 25, 29, 30]. Three examiners (Two endodontists and an oral radiologist with more than 15 years of experience) evaluated all the scans twice. The examiners were calibrated at the beginning of the study by evaluating 15% of the scans twice with two

weeks between the examinations. Any disagreement between the observers in the evaluation was resolved through discussion with a fourth observer (an endodontist with more than 10 years of experience).

Observed morphological features

Mandibular second molars were analysed for the following: (i) number of roots, (ii) presence and prevalence of supernumerary roots; radix entomolaris or radix paramolaris, (iii) number of root canals, (iv) prevalence of C-shaped root canal system configuration in which the examined molars should have the following features: (a) fused roots, (b) a longitudinal groove on lingual or buccal surface of the root, and (c) at least one cross-section of the canal belongs to the Fan et al. [18] criteria (Fig. 1), (v) correlation of prevalence of C-shaped root canal system configuration with gender, (vi) morphology of C-shaped molars according to Fan et al. [18]; (C1): a continuous C-shaped root canal with no separation or division, (C2): a comma-shaped root canal, resulting in a non-continuous C-shaped root canal outline (mesio-buccal-distal canal and a mesio-lingual canal), (C3a): two separate root canals, (C3b): three separate root canals, (C4): only one round, oval or flat canal in the cross-section, (C5): no presence of a canal, (vii) morphology of non-C-shaped mandibular molars according to Vertucci classification (Fig. 2) [32].

Statistical analysis

Weighted coefficient kappa (Kw) was used to measure interobserver reproducibility between observers separately for each time period and to measure intraobserver reproducibility between time periods separately for each observer. Categorical data were presented as frequency and percentage values and were analyzed using chi square test followed by pairwise comparisons utilizing multiple Fisher’s exact tests with Bonferroni correction. Confidence intervals were calculated using the Wilson score interval method [33]. The significance level was set at $p < 0.05$ within all tests. Statistical analysis was performed with R statistical analysis software version 4.1.3 for Windows.

Results

There was an excellent intraobserver (Kw value, 0.97) and interobserver (Kw value, 0.96) reproducibility.

Root morphology and number of roots

The initial analysis showed that 87.1% (95%CI 83.2%:90.3%) of the mandibular second molars were non-C-shaped while 12.9% (95%CI 9.7%:16.8%) were C-shaped, and this was non-significant in relation to gender ($\chi^2 = 0.19, p = 0.656$).

In terms of the number of roots, two roots was the most common (83.4%) (95%CI 79.2%:87.0%) followed by presence of a single root (16.3%) (95%CI 12.8%:20.5%), and only one sample (0.3%) (95%CI



Fig. 1 Representative axial CBCT sections showing the classification by Fan et al. [18]. 1a: class C1, a continuous C shaped canal. 1b: class C2, a mesiobuccal-distal (MB-D) ribbon-like canal and a mesiolingual (ML) canal. 1c: class C3a, a mesial (M) canal and a distal (D) canal. 1d: class C3b, a mesiobuccal (MB), a ML and a D canal. 1e: class C4, a single round or oval canal




Fig. 2 Root canal configurations detected in the study according to Vertucci’s classification (Types I to VI)

0.1%:1.6%) had three roots, and this was also non-significant in relation to gender ($\chi^2 = 1.86, p = 0.431$) (Table 1, Figs. 3, 4, 5 and 6).

Number of roots canals

Most samples had three (80%) (95%CI 75.5%:83.9%) followed by two (16%) (95%CI 12.5%:20.2%), then one

(3.1%) (95%CI 1.8%:5.5%), and finally four (0.9%) (95%CI 0.3%:2.5%) root canals for both genders (Table 2, Fig. 7). Yet, the probability of occurrence of three root canals was significantly higher in females and the probability of one or two root canals was significantly higher in males ($\chi^2 = 33.33, p < 0.001$).

Table 1 Number of roots and prevalence of C-shaped anatomy in relation to gender

Number of roots		Males	Females	Total	χ^2	p-value
One	n(%)	21(6%)	36(10.2%)	57(16.3%)	1.86	0.431
Two	n(%)	109(31.1%)	183(52.2%)	292(83.4%)		
Three (Radix Paramolaris)	n(%)	0(0.0%)	0(0.0%)	0(0.0%)		
Three (Radix Entomolaris)	n(%)	1(0.2%)	0(0.0%)	1(0.3%)		
Total	n(%)	131(37.4%)	219(62.6%)	350(100.0%)		
C-shaped		Males	Females	Total	χ^2	p-value
Yes	n(%)	16(4%)	29(8.2%)	45(12.9%)	0.19	0.656
No	n(%)	119(34.0%)	186(53.1%)	305(87.1%)		

$p = 0.431$ (non-significant) and $p = 0.656$ (non-significant)




Fig. 3 Pie chart showing number of roots and supernumerary roots



Fig. 4 The only sample of the study showing three roots (Radix entomolaris – 0.29%)




Fig. 5 Pie chart showing the percentage of C-shaped molars




Fig. 6 Samples of the C shaped molars included in the study

Table 2 Number of roots canals in relation to gender

		Males	Females	Total	χ_2	p-value
One	n(%)	8(2.2%)	3(0.9%)	11(3.1%)	33.33	< 0.001*
Two	n(%)	38(10.9%)	18(5.1%)	56(16%)		
Three	n(%)	87(24.9%)	193(55.1%)	280(80%)		
Four	n(%)	2(66.7%)	1(33.3%)	3(0.9%)		
Total	n(%)	135(38.6%)	215(61.4%)	350(100.0%)		

* significant ($p < 0.05$)




Fig. 7 Pie chart showing the percentage of molars showing a specific number of root canals

Canal morphology of C-shaped molars

There was a significant difference between different forms of the C-shaped molars within different root sections ($\chi^2 = 52.59, p < 0.001$) (Table 3). Post hoc pairwise comparisons showed the C1 configuration to be the most common in the coronal one third (44.4%) while C3b was the most prevalent in both the middle and apical thirds (28.9%) and (62.2%), respectively.

Root canals configuration of 2-rooted second molars according to Vertucci's classification

There was a significant difference between different root canal configurations in different roots ($\chi^2 = 417.72, p < 0.001$) (Table 4). Type IV Vertucci was the most common configuration found in the mesial roots while Type I was the most prevalent one in the distal roots.

Cross-sectional root configuration for non-C molars with 2 separate roots (n = 269)

For the mesial root, there was a significant difference between different root canal configurations in different sections ($\chi^2 = 74.59, p < 0.001$) (Table 5). The long oval configuration was the most commonly found in mesial roots while the oval cross section was the most prevalent

Table 3 Prevalence of C-shaped classes according to Fan et al. (18) in different root thirds

Section		C1 ^A	C2 ^{AB}	C3a ^{BC}	C3b ^C	C4 ^C	Total	χ_2	p-value
Coronal	n(%)	20(44.4%)	12(26.7%)	4(8.9%)	7(15.6%)	2(4.4%)	45(100.0%)	52.59	< 0.001*
Middle	n(%)	8(17.8%)	11(24.4%)	7(15.6%)	13(28.9%)	6(13.3%)	45(100.0%)		
Apical	n(%)	1(2.2%)	0(0.0%)	5(11.1%)	28(62.2%)	11(24.4%)	45(100.0%)		

Groups with different superscript letters within the heading row are significantly different; *significant ($p < 0.05$)

Table 4 Root canal configuration according to Vertucci's classification in mandibular second molars with 2 roots (n = 292)

Root		I ^A	II ^B	III ^B	IV ^C	V ^{BC}	VI ^{BC}	Total	χ_2	p-value
Mesial	n(%)	24(8.2%)	44(15.1%)	16(5.5%)	201(68.8%)	5(1.7%)	2(0.7%)	292(100.0%)	417.72	< 0.001*
Distal	n(%)	268(91.8%)	11(3.8%)	6(2.1%)	5(1.7%)	1(0.3%)	1(0.3%)	292(100.0%)		

Groups with different superscript letters within the heading row are significantly different; *significant ($p < 0.05$)

Table 5 Cross-sectional root configuration for non-C molars with 2 separate roots ($n = 269$)

Root	Section		Bowling pin ^A	Kidney shaped ^B	Long oval ^B	Oval ^B	Ribbon ^B	Round ^{AB}	Total	χ_2	p-value
Mesial	Coronal	n(%)	22(8.2%)	61(22.7%)	157(58.4%)	3(1.1%)	26(9.7%)	0(0.0%)	269(100.0%)	74.59	<0.001*
	Middle	n(%)	47(17.5%)	66(24.5%)	156(58.0%)	0(0.0%)	0(0.0%)	0(0.0%)	269(100.0%)		
	Apical	n(%)	22(8.2%)	65(24.2%)	182(67.7%)	0(0.0%)	0(0.0%)	0(0.0%)	269(100.0%)		
Root	Section		Bowling pin ^{ABC}	Kidney shaped ^A	Long oval ^B	Oval ^C	Ribbon ^{ABC}	Round ^B	Total	χ_2	p-value
Distal	Coronal	n(%)	2(0.7%)	68(25.3%)	37(13.8%)	101(37.5%)	2(0.7%)	59(21.9%)	269(100.0%)	219.54	<0.001*
	Middle	n(%)	0(0.0%)	84(31.2%)	0(0.0%)	181(67.3%)	0(0.0%)	4(1.5%)	269(100.0%)		
	Apical	n(%)	0(0.0%)	53(19.7%)	0(0.0%)	216(80.3%)	0(0.0%)	0(0.0%)	269(100.0%)		

Groups with different superscript letters within the same heading row are significantly different; *significant ($p < 0.05$)

one in distal roots; both in all three investigated levels of the root.

Discussion

The wide variation in radicular anatomy remains a major challenge in Endodontic practice [2]. These variations are not only limited to the number of root canals but also include the number of roots, root canals configuration and cross-sectional shapes [34]. These variations exist, not only between different ethnic populations, but even among individuals of the same ethnicity [35]. The Egyptian population is rich in diversity. By means of race it is considered Mediterranean, a sub-race of Caucasian, and populate the center of the Middle East-North Africa region. This study was done to address the knowledge gap related to the root and canal morphology of mandibular second molars in Egyptians.

CBCT scans were used in this investigation because of its credibility, wide acceptance and the large number of studies that employed CBCT in their methodology, thus allowing for direct comparisons [24–29]. Our inclusion criteria comprised only high-resolution, low field of view scans. This is a strength point for this study, because such imaging parameters provide better image quality and enhanced assessment of all investigated anatomic features. Another strength point is the number of experienced examiners ($n=3+1$) who assessed the scans twice. Although micro-CT can provide further descriptive anatomical details [36], it requires more scan time with high radiation doses and is limited to ex-vivo investigations. C-shaped anatomy configuration was studied according to Fan et al. [18]. This classification was chosen over the unmodified classification by Melton et al. [37] because the former was employed in numerous studies, hence allow the possibility of direct comparisons.

Results showed that the majority of mandibular second molars in Egyptians had two roots (83.4%), A third root was prevalent in only one sample (0.29%) of the studied subpopulation. This agrees similar studies in Caucasians (1.8%-2.7%) [26, 38], but disagrees others in Eskimos, North Americans, Indians and Koreans who had higher percentages (25.3%-51.4%) of a third root [39].

Regarding the number and descriptive morphology of root canals, most samples had three (80%) followed by two (16%), then one (3.2%), and finally four (0.8%) canals. This agrees a previous study in an Egyptian subpopulation [40], as well as similar findings in Koreans [12] and Venezuelans [41], also in Turks [42] except that they found the four-canals configuration to be slightly more than the single-canal configuration.

When the internal anatomy was analysed, the mesial root showed Vertucci type IV (68.8%) configuration to be the most prevalent one in Egyptians followed by Vertucci type II (15.1%). This agrees similar findings in Turks [42] and Iranians [43] but is opposite to findings reported in Yemenis [30] and Venezuelans [41]. For the distal root, the most prevalent configuration was Vertucci type I (91.8%), which comes in accordance with most similar studies [30, 40–43].

Root cross sections were also examined and classified. The most common cross sections were long oval in the mesial root and oval in the distal root followed by the kidney shape in both. This disagrees the findings of Senan et al. [30] in Yemenis where the mesial root was mostly ribbon-shaped and the distal root was kidney-shaped. Radicular cross sections need to be studied on case-by-case basis to be able to recognize when the dentin thickness poses a threat in the dangerous zone [44].

The C-shaped molars in this study were found in 12.8% of the examined scans. This percentage is close to that reported in Belgians (10.7%) [6], Sudanese (10%) [45], Saudis (9.1%) [46], Yemenis (9%) [30], Chileans (8.9%) [6], Indians (9.7%, 8.1%) [47, 48], Brazilians (8.5%) [49] and Lebanese (9.09%) [50]. However, it was lower than results in Iranians (17.6%, 21.4%) [43, 51], Chinese (29%, 38.6%, 47.05%) [4, 52, 53], Koreans (39.8%, 44.5%) [54, 55] and Malaysians (48.7%) [56]. Nonetheless, it was higher than results in Turks (4.1%) [42]. This 12.8% prevalence of C-shaped canals contrasts another study done in an Egyptian subpopulation [40]. This contrast may be attributed to the differences in sample size, and more importantly, the imaging parameters.

The C-shaped molars were found more in female than male Egyptians with no significant difference between

them. This agrees with reports from Chinese [4], Venezuelans [41], Indians [48] and Iranians [43]. However, significant gender differences was reported by Sharaan and Elrawdy in another Egyptian subpopulation [40] and was found in Brazilians [49], Koreans [55], Saudis [46], Malaysians [56], Portuguese [13], and Turkish [16] populations with a higher incidence of C-shaped molars among females. Such contrasts emphasize the existing inter- and intra-ethnic variations and highlight the importance of CBCT as a clinical adjunct when managing individual cases.

Regarding canal configuration of C-shaped molars, C1 was the prevalent coronal configuration, while C3b was the most prevalent one in both the middle and apical thirds. This totally agrees reports in Chinese [4], Venezuelan [41] and Iranian [43] populations, partially agrees reports in Portuguese [13], Turkish [16], and Yemini [30] populations who described different middle and apical configurations. However, disagree Alfawaz et al. [46] who found C3a to be the most prevalent coronal configuration, and C3b to be the most prevalent in middle and apical configurations in Saudis, and Kim et al. [55] who reported C2 to be the most prevalent coronal configuration in Koreans.

It is worth mentioning that this classification of Fan et al. (18) was misquoted by some studies as they used the designations C3c and C3d instead of C3a and C3b respectively [4, 30, 46, 57]. This was probably caused by the confusing labels of the demonstrative figure in Fan et al. [18] that was republished later with clearer labeling [19]. Though this remark is not exactly a finding, it is still worth mentioning as a reminder of the importance of revising primary references and providing clear illustrations.

This study adds to the scarce knowledge available about the studied population [40]. Also, the study has significant clinical relevance; Studies have shown that fused roots molars have a higher association with periapical lesions when compared to non-fused teeth [58]. Thus, the study findings that show the wide range of morphological variations justify a preoperative limited field of view CBCT for mandibular second molars once endodontic treatment is indicated. This agrees the AAE/AAOMR joint statement [59].

This study has the limitation of not investigating the ethnicity of the patients because this information was not available to collect and analyze. Also, the symmetric distribution of findings was not possible because only small field of view scans were included for the anatomic analysis. Nonetheless, even studies that showed a high probability of a patient to have C-shaped canals bilaterally still recommend individual evaluation of each mandibular second molar while performing endodontic treatment

on both sides [4, 30, 40]. Finally, the impact of age, which can affect the root canal anatomy, was not included in the analysis because they were a minority in the studied population. In which about 86.6% of the patients were in the age range of 21–50 years.

In conclusion, Egyptian permanent mandibular second molars have mainly two separated mesial and distal roots with a very low prevalence of a third one. Various canals configurations were identified, with prevalence for Vertucci Type IV and Type I in the mesial and distal roots, respectively. The occurrence of C-shaped roots and canals must be considered when treating these molars, mainly in females. Also, high resolution, low field of view CBCT is a useful clinical adjunct for successful management of such teeth.

Abbreviations

CBCT	Cone beam computed tomography
FOV	Field of view
ICC	Interclass correlation coefficient

Acknowledgements

N/A.

Authors' contributions

All authors (S.S., M.A.S, S.A.E.S and N.N.N) contributed to the conception, design and execution of the study. S.S. and N.N.N drafted the work and substantively revised it. All authors have viewed and approved the initially submitted version (and the revised ones) and have agreed to be personally accountable to the submitted work.

Funding

Open access funding provided by The Science, Technology & Innovation Funding Authority (STDF) in cooperation with The Egyptian Knowledge Bank (EKB). Self-funded.

Availability of data and materials

The data that support the findings of this study are available from The British University—faculty of Dentistry, but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the authors upon reasonable request and with permission of The British University—faculty of Dentistry.

Declarations

Ethics approval and consent to participate

This study was approved and the need for participation consents was waived by the Research Ethics Committee of The British University (El-Shorouk city, Egypt) (22–007). All procedures performed were in accordance with the 1964 Helsinki declaration and its later amendments.

Consent for publication

N/A.

Competing interests

The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Author details

¹Department of Endodontics, Faculty of Dentistry, The British University in Egypt (BUE), El Sherouk City, Cairo, Egypt. ²Center for Innovative Dental Sciences (CIDS), Faculty of Dentistry, The British University in Egypt (BUE), Cairo, Egypt. ³Department of Endodontics, Faculty of Dentistry, Ain Shams University,

Cairo, Egypt. ⁴Department of Oral and Maxillofacial Radiology, Ain Shams University, Cairo, Egypt.

Received: 24 June 2022 Accepted: 4 April 2023

Published online: 15 April 2023

References

- Vertucci F. Root canal morphology and its relationship to endodontic procedures. *Endod Top*. 2005;10:3–29.
- Cantatore G, Berutti E, Castellucci A. Missed anatomy: frequency and clinical impact. *Endod Top*. 2006;15:3–31.
- von Zuben M, Martins JNR, Berti L, Cassim I, Flynn D, Gonzalez JA, Gu Y, Kottoor J, Monroe A, Rosas Aguilar R, Marques MS, Ginjeira A. Worldwide Prevalence of Mandibular Second Molar C-Shaped Morphologies Evaluated by Cone-Beam Computed Tomography. *J Endod*. 2017;43:1442–7.
- Zheng Q, Zhang L, Zhou X, Wang Q, Wang Y, Tang L, Song F, Huang D. C-shaped root canal system in mandibular second molars in a Chinese population evaluated by cone-beam computed tomography. *Int Endod J*. 2011;44:857–62.
- Gulabivala K, Aung TH, Alavi A, Ng YL. Root and canal morphology of Burmese mandibular molars. *Int Endod J*. 2001;34:359–70.
- Torres A, Jacobs R, Lambrechts P, Brizuela C, Cabrera C, Concha G, Pedemonte ME. Characterization of mandibular molar root and canal morphology using cone beam computed tomography and its variability in Belgian and Chilean population samples. *Imaging Sci Dent*. 2015;45:95–101.
- Gulabivala K, Opasanon A, Ng YL, Alavi A. Root and canal morphology of Thai mandibular molars. *Int Endod J*. 2002;35:56–62.
- Al-Qudah AA, Awawdeh LA. Root and canal morphology of mandibular first and second molar teeth in a Jordanian population. *Int Endod J*. 2009;42:775–84.
- Peiris R. Root and canal morphology of human permanent teeth in a Sri Lankan and Japanese population. *Anthropol Sci*. 2008;116:123–33.
- Al ME, Chaudhry J, Abdo S, Rashid EA, Khamis AH, Jamal M. Evaluation of root and canal morphology of maxillary permanent first molars in an Emirati population; a cone beam computed tomography study. *BMC Oral Health*. 2020;20:274–274.
- Fayyaz A, Patil S, Maragathavalli G, Kumar R. Prevalence and configurations of C-shaped canals in permanent mandibular second molars in a Saudi Arabian population: A CBCT study. *Int Med J*. 2019;26:223–5.
- Kim SY, Kim BS, Kim Y. Mandibular second molar root canal morphology and variants in a Korean subpopulation. *Int J Endod*. 2016;49:136–44.
- Martins JNR, Mata A, Marques D, Carames J. Prevalence of C-shaped mandibular molars in the Portuguese population evaluated by cone-beam computed tomography. *Eur J Dent*. 2016;10:529–35.
- Ladeira DB, Cruz AD, Freitas DQ, Almeida SM. Prevalence of C-shaped root canal in a Brazilian subpopulation: A cone-beam computed tomography analysis. *Braz Oral Res*. 2014;28:39–45.
- Silva EJ, Nejaim Y, Silva AV, Haiter-Neto F, Cohenca N. Evaluation of root canal configuration of mandibular molars in a Brazilian population by using cone-beam computed tomography: an in vivo study. *J Endod*. 2013;39:849–52.
- Kaplan SS, Kaplan T, Sezgin GP. Evaluation of C-shaped canals in mandibular second molars of a selected patient group using cone beam computed tomography: prevalence, configuration and radicular groove types. *Odontology*. 2021;109:949–55.
- Cooke HG, Cox FL. C-shaped canal configurations in mandibular molars. *J Am Dent Assoc*. 1979;99:836–9.
- Fan B, Cheung GSP, Fan M, Gutmann JL, Bian Z. C-shaped canal system in mandibular second molars: part I – anatomical features. *J Endod*. 2004;30:899–903.
- Fan W, Fan B, Gutmann JL, Cheung GS. Identification of C-shaped canal in mandibular second molars. Part I: radiographic and anatomical features revealed by intraradicular contrast medium. *J Endod*. 2007;33:806–10.
- Chai WL, Thong YL. Cross-sectional morphology and minimum canal wall widths in C-shaped roots of mandibular molars. *J Endod*. 2004;30:509–12.
- Fernandes M, de Ataíde I, Wagle R. C-shaped root canal configuration: a review of literature. *J Conserv Dent*. 2014;17:312–9.
- Calberson P, De Moor R, Deroose C. The radix entomolaris and paramolaris: clinical approach. *J Endod*. 2007;33:58–63.
- Ahmed HMA. A critical analysis of laboratory and clinical research methods to study root and canal anatomy. *Int J Endod*. 2022;55:229–80.
- Patel S, Durack C, Abella F, Shemesh H, Roig M, Lemberg. Cone beam computed tomography in Endodontics-a review. *Int J Endod*. 2015;48:3–15.
- Saber S, El Sadat SA, Taha A, Nawar NN, Azim AA. Anatomical Analysis of Mandibular Posterior Teeth using CBCT: An Endo-Surgical Perspective. *Eur Endod J*. 2021;6:264–70.
- Martins JN, Marques D, Mata A, Carames J. Root and root canal morphology of the permanent dentition in a Caucasian population: a cone-beam computed tomography study. *Int J Endod*. 2017;50:1013–26.
- Alemam S, Abuelsadat S, Shehabeldin S, Elsewif T. Accuracy, sensitivity and specificity of three imaging modalities in detection of separated intracanal instruments. *Italian Endod J*. 2020;34:97–103.
- Ghobashy AM, Nagy MM, Bayoumi AA. Evaluation of root and canal morphology of maxillary permanent molars in an Egyptian population by cone-beam computed tomography. *J Endod*. 2017;43:1089–92.
- Saber SEDM, Ahmed MHM, Obeid M, Ahmed HMA. Root and canal morphology of maxillary premolar teeth in an Egyptian subpopulation using two classification systems: a cone beam computed tomography study. *Int J Endod*. 2019;52:267–78.
- Senan EM, Alhadainy HA, Madfa AA. Root and canal morphology of mandibular second molars in a Yemeni population: A cone-beam computed tomography. *Eur Endod J*. 2021;6:72–81.
- Martins JNR, Kishen A, Marques D, et al. Preferred reporting items for epidemiologic cross-sectional studies on root and root canal anatomy using cone-beam computed tomographic technology: A systematized assessment. *J Endod*. 2020;46:915–35.
- Vertucci FJ. Root canal anatomy of the human permanent teeth. *Oral Surg Oral Med Oral Pathol*. 1984;58:589–99.
- Wilson EB. Probable inference, the law of succession, and statistical inference. *JASA*. 1927;22:209–12.
- Razumova S, Brago A, Howijieh A, Barakat H, Kozlova Y, Baykulova M. Evaluation of Cross-Sectional Root Canal Shape and Presentation of New Classification of Its Changes Using Cone-Beam Computed Tomography Scanning. *Appl Sci*. 2020;10:4495.
- Razumova S, Brago A, Khaskhanova L, Howijieh A, Barakat H, Manvelyan A. A Cone-Beam Computed Tomography Scanning of the Root Canal System of Permanent Teeth among the Moscow Population. *Int J Dent*. 2018;2018:2615746. <https://doi.org/10.1155/2018/2615746>.
- Elashiry M, Shehab Eldin S, Elashry S. Comparison of Shaping Ability of Different Single-File Systems Using Microcomputed Tomography. *Euro J Dent*. 2020;14:70–6.
- Melton DC, Krell KV, Fuller MW. Anatomical and histological features of C-shaped canals in mandibular second molars. *J Endod*. 1991;17:384–8.
- Ferraz JA, Pécora JD. Three-rooted mandibular molars in patients of Mongolian, Caucasian and Negro origin. *Braz Dent J*. 1993;3:113–7.
- Kantilieraki E, Delantoni A, Angelopoulos C, Beltes P. Evaluation of Root and Root Canal Morphology of Mandibular First and Second Molars in a Greek Population: A CBCT Study. *Eur Endod J*. 2019;4:62–8.
- Sharaan ME, Elrawdy AM. An evaluation of mandibular molars root canal morphology using cone-beam computed tomography in an Egyptian subpopulation. *Tanta Dent J*. 2017;14:220–4.
- Gomez F, Brea G, Gomez-Sosa JF. Root canal morphology and variations in mandibular second molars: an in vivo cone-beam computed tomography analysis. *BMC Oral Health*. 2021;21:424.
- Demirbuga S, Sekerci AE, Dinçer AN, Cayabatmaz M, Zorba YO. Use of cone-beam computed tomography to evaluate root and canal morphology of mandibular first and second molars in Turkish individuals. *Med Oral Patol Oral Cir Bucal*. 2013;18:737–44.
- Janani M, Rahimi S, Jafari F, Johari M, Nikniaz S, Ghasemi N. Anatomic Features of C-shaped Mandibular Second Molars in a Selected Iranian Population Using CBCT. *Iran Endod J*. 2018;13:120–5.
- Sauáia TS, Gomes BPFA, Pinheiro ET, Zaia AA, Ferraz CCR, Souza-Filho FJ, Valdrighi L. Thickness of dentine in mesial roots of mandibular molars with different lengths. *Int J Endod*. 2010;43:555–9.
- Ahmed HA, Abu-bakr NH, Yahia NA, Ibrahim YE. Root and canal morphology of permanent mandibular molars in a Sudanese population. *Int J Endod*. 2007;40:766–71.

46. Alfawaz H, Alqedairi A, Alkhayyal AK, et al. Prevalence of C-shaped canal system in mandibular first and second molars in a Saudi population assessed via cone beam computed tomography: a retrospective study. *Clin Oral Invest.* 2019;23:107–12.
47. Felsypremila G, Vinothkumar TS, Kandaswamy D. Anatomic symmetry of root and root canal morphology of posterior teeth in Indian subpopulation using cone beam computed tomography: A retrospective study. *Eur J Dent.* 2015;9:500–7.
48. Wadhvani S, Singh MP, Agarwal M, Somasundaram P, Rawtiya M, Wadhvani PK. Prevalence of C-shaped canals in mandibular second and third molars in a central India population: A cone beam computed tomography analysis. *J Conserv Dent.* 2017;20:351–4.
49. Nejaim Y, Gomes AF, Rosado LPL, Freitas DQ, Martins JNR, da Silva EJNL. C-shaped canals in mandibular molars of a Brazilian subpopulation: prevalence and root canal configuration using cone-beam computed tomography. *Clin Oral Invest.* 2020;24:3299–305.
50. Feghali M, Jabre C, Haddad G. Anatomical Investigation of C-shaped Root canal systems of mandibular molars in a middle-eastern population: A CBCT Study. *J Contemp Dent Pract.* 2022;23:713–9.
51. Madani ZS, Mehraban N, Moudi E, Bijani A. Root and canal morphology of mandibular molars in a selected Iranian population using cone-beam computed tomography. *Iran Endod J.* 2017;12:143–8.
52. Zhang R, Wang H, Tian YY, Yu X, Hu T, Dummer PM. Use of cone-beam computed tomography to evaluate root and canal morphology of mandibular molars in Chinese individuals. *Int Endod J.* 2011;44:990–9.
53. Chen C, Zhu T, Wu H, et al. Prevalence and correlation of C-shaped root canals of mandibular premolars and molars in Eastern Chinese individuals. *Sci Rep.* 2022;12:19779.
54. Jin GC, Lee SJ, Roh BD. Anatomical study of C-shaped canals in mandibular second molars by analysis of computed tomography. *J Endod.* 2006;32:10–3.
55. Kim HS, Jung D, Lee H, Han YS, Oh S, Sim HY. C-shaped root canals of mandibular second molars in a Korean population: a CBCT analysis. *Restor Dent Endod.* 2018;43:e42.
56. Pan JYY, Parolia A, Chuah SR, Bhatia S, Mutalik S, Pau A. Root canal morphology of permanent teeth in a Malaysian subpopulation using conebeam computed tomography. *BMC Oral Health.* 2019;19:14.
57. Jafarzadeh H, Wu YN. The C-shaped root canal configuration: a review. *J Endod.* 2007;33:517–23.
58. Pereira B, Martins JNR, Baruwa AO, et al. Association between endodontically treated maxillary and mandibular molars with fused roots and periapical lesions: A cone-beam computed tomography cross-sectional study. *J Endod.* 2020;46:771–7.
59. Special committee to revise the joint AAE/AAOMR position statement on use of CBCT in endodontics. AAE and AAOMR joint position statement: Use of cone beam computed tomography in endodontics 201 update. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2015;120:508–12.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

